

L3(3.3)-Quadratic Relations in Factored Form

## Key Concepts:

- factored form of quadratic relation
- direction of opening from 'a'
- solving for zeroes

- using symmetry to find:
  - x-coordinate of vertex
  - axis of symmetry

- using substitution to find:
  - y-coordinate of vertex
  - y-intercept

Is  $y = 2(x+1)(x-5)$  a quadratic relation?

Examine 1st and 2nd differences:

x	y	1st	2nd
-2	14		
-1	0	$0 - (14) = -14$	$-10 - (-14) = 4$
0	-10	$-10 - (0) = -10$	$-6 - (-10) = 4$
1	-16	$-16 - (-10) = -6$	$-2 - (-6) = 4$
2	-18	$-18 - (-16) = -2$	

Quadratic

Quadratic Relations in Factored Form

The equation of a quadratic relation may be written in several forms:

1. standard form:  $y = ax^2 + bx + c$

2. factored form:  $y = a(x - s)(x - t)$

3. vertex form:  $y = a(x - h)^2 + k$

*method*

The factored form,  $y = a(x - s)(x - t)$ , is most useful for finding the zeroes, which are  $x = s$  and  $x = t$ .

Consider the following...

Give two numbers that have a product of zero:

★  $0(10) = 0$        $(0)(0) = 0$   
 $10(0) = 0$        $(5+5) = 0$

What do you notice? (any value)  $\times 0 = 0$

★ only way to get a product of zero is  $x \times 0$


Solve:

(a)  $\frac{3x}{3} = \frac{0}{3}$   
 $x = 0$

(b)  $\frac{57y}{57} = \frac{0}{57}$   
 $y = 0$

(c)  $\frac{3xy}{3} = \frac{0}{3}$   
 $xy = 0$

Depending upon the location of the vertex, and whether the parabola opens up or down, it may have 0, 1, or 2 distinct (unique) zeroes.



Zeroes occur where the y-coordinate of the parabola is equal to zero.

To find the zeroes algebraically, we set  $y = 0$  and solve for the x-values that make the equation true.

Ex.1 Determine the zero(es) of each

(a)  $y = x(x - 10)$

$$0 = x(x - 10)$$

$x = 0$  or  $x = 10$

Recall:

Zero multiplied by anything is zero.

If  $(a)(b) = 0$  then  
 $a = 0$  or  $b = 0$  (or both are zero).

(b)  $y = -2(x - 5)(3x - 1)$

$$0 = -2(x - 5)(3x - 1)$$

$x = 5$  or  $3x - 1 = 0$

$$\frac{3x}{3} = \frac{1}{3}$$

$$x = \frac{1}{3}$$

2 Zeroes

(c)  $y = 2(x - 2)^2$

$$0 = 2(x - 2)(x - 2)$$

$x = 2$  or  $x = 2$

1 Zero

The zeroes and symmetry can be used to find the vertex (h, k).

For the x-coordinate (h), find the midpoint of the zeroes:

$$\star \quad MP = \frac{x_1 + x_2}{2} = \frac{s + t}{2}$$

For the y-coordinate (k), substitute the midpoint into the equation and solve for y:

$$\star \quad y = a(x - s)(x - t)$$

$$y = a(MP - s)(MP - t)$$

Steps

1. find zeroes

2. find axis of symmetry

3. Sub AOS into equation to solve k

Ex.2 Determine the vertex:

(a)  $y = -2(x - 2)(x - 8)$

$$0 = -2(x - 2)(x - 8)$$

$$x = 2 \text{ or } x = 8$$

$$AOS = \frac{2 + 8}{2}$$

$$x = 5$$

Sub  $x = 5$  into equation

$$= -2(5 - 2)(5 - 8)$$

$$= -2(3)(-3)$$

$$= 18$$

$\therefore$  the vertex is (5, 18)

Ex.3 A parabola has zeroes at -3 and 2, and a y-intercept of 18. Determine the equation.

$$y = a(x - s)(x - t)$$

$$y = a(x - (-3))(x - (2))$$

$$y = a(x + 3)(x - 2)$$

y-intercept (0, 18)

$$18 = a(0 + 3)(0 - 2)$$

$$18 = a(3)(-2)$$

$$18 = \frac{-6a}{-6}$$

$$a = -3$$

$\therefore$  the equation is

$$y = -3(x + 3)(x - 2)$$

Assigned Work: p. 155-157 # 2, 3, 4ace, 5, 6ace, 7, 10

